

RESEARCH ARTICLE

Naser Maheri-Sis¹
Mehdi Eghbali-Vaighan¹
Ali Mirzaaghazadeh²
Yahya Ebrahimnezhad¹

Effect of microwave irradiation on the fermentation characteristics and nutritive value of tomato pomace for ruminants using *in vitro* gas production technique

Authors' addresses:

¹Department of Animal Science, Shabestar Branch, Islamic Azad University, 5381637181 Shabestar, East Azerbaijan, Iran.

²Department of Animal Science, Urmia University, Urmia, Iran.

Correspondence:

Naser Maheri-Sis
Department of Animal Science, Shabestar Branch, Islamic Azad University, 5381637181 Shabestar, East Azerbaijan, Iran.
Tel.: +98-471-2225311
e-mail: maherisis@iaushab.ac.ir

Article info:

Received: 7 November 2012

Accepted: 28 December 2012

ABSTRACT

This study was conducted to estimate the effect of microwave irradiation on fermentation pattern, metabolizable energy content and organic matter digestibility of tomato pomace using *in vitro* gas production technique. Tomato pomace samples were collected, pooled, ground and then exposed to 1000W microwave irradiation for 0, 2.5 and 5 minutes. The experimental samples were incubated *in vitro* with rumen liquor taken from three fistulated Iranian native cows at 2, 4, 6, 8, 12, 16, 24, 36, 48, 72 and 96 h. Cumulative gas production volume at early incubation times was significantly decreased by microwave treating time ($P < 0.01$). Microwave irradiation could not significantly affect gas production parameters. Significant effects were not found on metabolizable energy content (7.82 - 8.02 MJ/ Kg DM) and organic matter digestibility (51.87% - 53.15%) of microwave treated tomato pomace. The results showed that microwave irradiation may be a useful method to decrease ruminal gas production (including methane), without any negative effects on nutritional value of tomato pomace.

Key words: tomato pomace, microwave, nutritive value, gas production, metabolizable energy

Introduction

Agro-industrial by-products refer to the by-products derived in the industry due to processing of main products. Many of these agro-industrial by-products have a substantial potential nutritional value as livestock feeds. Thus, feeding such by-products will help to decrease feeding cost especially in developing countries (Sindhu et al., 2002). The ruminant animals have the unique capacity to utilize fibrous by-products because of their rumen microbes. Ruminal bacteria, protozoa and fungi together contribute to microbial fermentation of plant cell walls in the rumen (Chaji et al., 2011). Using agro-industrial by-products in ruminant nutrition is one of the economic strategies to overtop feed shortage worldwide (Mirzaei-Aghsaghali & Maheri-Sis, 2008; Chaji et al., 2011; Taher-Maddah et al. 2012). Tomato pomace is an agro-industrial by-product which is producing in significant amounts in Iran. Tomato pomace produced in Iran contains 21.7-26.4% crude protein, 13.4-15.9% fat, 3.4-

12.2% crude ash and 49.2-57.4% neutral detergent fiber on dry matter basis and 26% DM on as-fed basis (Abdollahzadeh et al., 2010; Aghajanzadeh-Golshani et al., 2010; Mirzaei-Aghsaghali et al., 2011). Based on our previous studies nutritive value of dried tomato pomace can be comparable with alfalfa hay in ruminant nutrition (Maheri-Sis et al., 2007; Mirzaei-Aghsaghali et al., 2008; Aghajanzadeh-Golshani et al., 2010; Mirzaei-Aghsaghali et al., 2011).

Because of quick spoilage of wet tomato pomace, usually it is dried or ensiled for further using. Microwave irradiation is one of the most effective food and feed drying and processing methods which have been used in recent years (Sadeghi & Shawrang, 2008; Pelletier et al., 2010; Maheri-Sis et al., 2011a). Microwaves are non-ionizing electromagnetic waves positioned between the X-ray and infrared rays in the electromagnetic spectrum with frequency between 300 MHz to 300 GHz (Tatke & Jaiswal, 2011). Zhao et al., (2007) reviewed that microwave technology is

RESEARCH ARTICLE

increasingly playing an important role in drying in the food industry because of its rapid heating rate and ease of use. It also shows considerable potential for preventing mildew in food. In addition, this method of processing does not cause environmental pollution or introduce foreign chemical reagents. Chavan & Chavan (2010) suggested that microwave heating techniques have many unique advantages when compared to conventional heating methods because they are; convenient to operate and control, energy-efficient and clean. Other advantages of microwave irradiation include the following: shorter drying time, improved product quality and flexibility in producing a wide variety of dried products. Microwave irradiation has been used recently by Al-Harashsheha et al. (2009) for drying tomato pomace. They were found that drying rate have been increased with increasing microwave dosage. Aflatoxin contamination is one of the serious problems on conventional drying of tomato pomace. Farag et al. (1996) illustrated that microwave heating reduced the aflatoxin content considerably in contaminated material.

Effect of microwave irradiation on ruminal degradation of certain feedstuffs for ruminants has been studied recently by some researchers (Sadeghi & Shawrang, 2007; Ebrahimi et al., 2010, Maheri-Sis et al., 2011b, 2011c), but it seems that, there is needs to further investigations for understanding fermentation pattern and nutritive value of individual feeds and by-products.

Among three common techniques (*in vivo*, *in situ* and *in vitro*) used for nutritional evaluation of feeds, *in vitro* gas production technique is the superior method for feed evaluation (Ajmal Khan et al., 2003), especially in developing countries because it is capable of measuring rate and extent of nutrients fermentation with less expenditure and estimation of organic matter digestibility and metabolizable energy content of several classes of feedstuffs for ruminants (Menke et al., 1979; Maheri-Sis et al., 2008; Chen et al., 2011).

The aim of this study was to estimate the effect of microwave irradiation on fermentation pattern, metabolizable energy content and organic matter digestibility of tomato pomace using *in vitro* gas production technique.

Materials and Methods

Animals and feeds

Three fistulated Iranian native (Taleshi) cows fed twice daily with a diet containing hay (60%) and concentrate (40%) following the method described by Menke & Steingass

(1988), were used for rumen liquor collection in order to conduct in gas production technique. Tomato pomace samples were collected from tomato processing factories of Shabestar region, Iran. Three samples (500 g) were dried, pooled, ground (milled through a 1 mm sieve) and exposed to 1000W microwave irradiation (emitting a 2450 MHz microwave frequency) for 0, 2.5 and 5 minutes.

Chemical analysis

Tomato pomace samples milled through a 1 mm sieve for chemical analysis and gas production procedure. Dry matter (DM) was determined by drying the samples at 105°C overnight and ash by igniting the sample in muffle furnace at 550°C for 8 h. Nitrogen (N) content was measured by the Kjeldal method. Crude protein (CP) was calculated as $N \times 6.25$ (AOAC, 1990). Neutral detergent fiber (NDF) and acid detergent fiber (ADF) were determined by procedures outlined by Van Soest et al. (1991). Chemical analyses were carried out in animal nutrition laboratories of Islamic Azad University, Shabestar Branch and Animal Science Research Institute, Karaj, Iran.

In vitro gas production

The amount of 200 mg samples was weighed in triplicate into calibrated glass syringes of 100 ml. The syringes were prewarmed at 39°C before the injection of 30 ml rumen fluid-buffer mixture into each syringe followed by incubation in a water bath at 39°C. Readings of gas production were recorded before incubation (0) and 2, 4, 6, 8, 12, 16, 24, 36, 48, 72 and 96 h after incubation. Total gas values were corrected for blank incubation and gas production from syringes contain rumen fluid. Gas production procedure was done in laboratory of Animal Science Research Institute, Karaj, Iran.

Cumulative gas production data were fitted to the model of Ørskov & McDonald (1979):

$$Y = a + b(1 - e^{-ct})$$

where a = gas production from the immediately soluble fraction (mL), b = gas production from the insoluble fraction (mL), c = gas production rate constant for the insoluble fraction (h), $a + b$ = potential gas production (mL), t = incubation time (h) and Y = gas production at time t .

The metabolizable energy contents of tomato pomace were calculated using equations of Menke & Steingass (1988) as follows:

$$ME \text{ (MJ/kg DM)} = 2.20 + 0.136 \text{ GP} + 0.057 \text{ CP} + 0.0029 \text{ EE}^2$$

RESEARCH ARTICLE

where ME = metabolizable energy (MJ per kg DM), GP = 24 h net gas production (ml 200 mg⁻¹), CP = crude protein (%), EE = ether extract (%).

Organic mater digestibility of tomato pomace was calculated using equations of Menke *et al.* (1979) as below:

$$\text{OMD (\%)} = 0.9991 \text{ GP} + 0.595 \text{ CP} + 0.181 \text{ XA} - 9$$

where OMD = organic mater digestibility, GP = 24 h net gas production (ml 200 mg⁻¹), CP = crude protein (%), XA = ash content (%).

Statistical analysis

All data were analyzed based on completely randomized design (CRD) by using software of SAS (1991) and means (obtained from three samples) were separated by Duncan's multiple range tests (Steel & Torrie, 1980).

Results and Discussion

Chemical composition of tomato pomace was presented in Table 1. There are some differences and variations between chemical compositions in current study comparing with some other researches (Chumpawadee *et al.*, 2007; Besharati *et al.*, 2008; Chumpawadee, 2009; Aghajanzadeh-Golshani *et al.*, 2010; Mirzaei-Aghsaghali *et al.*, 2011). These

variations in chemical composition of by-products can be due to different original materials, growing conditions (geographic, seasonal variations, climatic conditions and soil characteristics), extent of foreign materials, impurities, tomato varieties, skin to seed ratio and different processing and measuring methods. It is predictable that, different chemical composition led to different nutritive value, because chemical composition is the initial and important index of nutritive value of feeds (Aghajanzadeh-Golshani *et al.*, 2010; Maheri-Sis *et al.*, 2011b, 2011c).

Table 1. Chemical composition of tomato pomace on dry matter basis (%).

DM	CP	EE	CA	ADF	NDF	NFC
94.46	24.08	13.98	5.73	33.6	47.8	8.41

Legend: DM - dry matter, CP - crude protein, EE - ether extract, CA - crude ash, ADF - acid detergent fiber, NDF - neutral detergent fiber, NFC - non fiber carbohydrate.

Gas production volume (ml / 200 mg DM) at different incubation times and gas production parameters (*a*, *b*, *a+b* and *c*) and calculated amounts of organic matter digestibility and metabolizable energy of untreated and microwave treated tomato pomace are presented in Figure 1 and Tables 2 and 3.




Figure 1. *In vitro* gas production volume of untreated and microwave treated (MT) tomato pomace at different incubation times.

RESEARCH ARTICLE

Table 2. Gas production volume (ml / 200mg DM) of microwave treated and untreated tomato pomace at different incubation times (h).

Incubation time (hours)	Untreated	Microwave treated (2.5 min)	Microwave treated (5 min)	P-value	S.E.M
2	6.67	6.35	6.06	0.1673	0.195
4	12.56 ^a	11.78 ^a	10.55 ^b	0.0040	0.253
6	18.60 ^a	17.03 ^b	15.04 ^c	0.0002	0.258
8	25.34 ^a	23.47 ^b	20.71 ^c	0.0003	0.359
12	36.34 ^a	35.40 ^a	33.30 ^b	0.0027	0.359
24	44.03	42.78	42.76	0.1686	0.468
48	46.23	44.35	45.43	0.1976	0.643
72	46.78	45.05	46.14	0.1899	0.585
96	47.49	45.89	46.85	0.2585	0.615

Means in the same row with different letters (a, b and c) differ (P<0.05).

Table 3. Gas production parameters, estimated metabolizable energy (ME) contents and organic matter digestibility (OMD) of untreated and microwave treated tomato pomace.

Items	Untreated	Microwave treated (2.5 min)	Microwave treated (5 min)	P-value	S.E.M
a (ml)	5.73	5.84	5.09	0.0963	0.236
b (ml)	52.89	51.39	51.83	0.1176	0.615
a+b (ml)	58.62	57.23	56.92	0.1284	0.325
c (ml.h ⁻¹)	0.114	0.113	0.096	0.0851	0.009
OMD (%)	53.15	51.89	51.87	0.4574	0.073
ME (MJ.Kg ⁻¹ DM)	8.02	7.82	7.82	0.2114	0.468

Legend: a - gas production from the immediately soluble fraction; b - gas production from the insoluble fraction; a+b - potential gas production; c - gas production rate constant for the insoluble fraction (b).

Cumulative gas production volume at early incubation times (until 12 h) was significantly decreased by microwave treating time (P<0.01). There was no significant difference between gas production parameters of untreated and microwave treated tomato pomace. The gas volume in all of incubation times and gas production parameters (a, b, a+b and c) for untreated tomato pomace were slightly higher than that reported by Besharati et al. (2008), Aghajanzadeh-Golshani et al. (2010) and Mirzaei-Aghsaghali et al. (2011). In our previous studies (Maheri-Sis et al., 2011b, 2011c) we found that microwave irradiation could not affect dry matter and protein degradability of tomato pomace. Karn (1991) previously had been reported that microwave drying have a minimal and inconsistent effect on organic matter digestibility of forages. Eskicioglu et al. (2007) stated that, neither the chemical mechanism of microwave interaction with materials, nor microbial destruction (sterilization)

mechanism of microwave in biological systems is fully understood. Maheri-Sis et al. (2011a) observed that microwave treatment lead to increasing *in vitro* gas production parameters of sunflower meal. It seems that nature of feedstuff could have a determinative role on efficiency of microwave irradiation.

It was confirmed that gas volume at 24 h after incubation correlate with the metabolizable energy in feedstuffs (Menke et al., 1979; Menke & Steingass, 1988; Chen et al., 2011). Additionally, *in vitro* organic matter digestibility was shown to have high correlation with gas production rate and volume (Sommart et al., 2000; Kilic & Garipoglu, 2009). In this study, microwave irradiation could not significantly affect metabolizable energy content (7.82 - 8.02 MJ/kg DM) and organic matter digestibility (51.87% - 53.15%) of tomato pomace. These values were lower than those of Besharati et al. (2008), Aghajanzadeh-Golshani et al. (2010) and Mirzaei-

RESEARCH ARTICLE

Aghsaghali et al. (2011), and approximately 50-60% higher than that of Chumpawadee et al. (2007). Different nutritive values obtained from various studies may be due to different chemical compositions (especially soluble carbohydrates, crude protein, non-fibrous carbohydrates, fat, acid detergent fibre and neutral detergent fibre), processing methods, tomato varieties, skin to seed ratio, climate conditions, evaluation procedures and experimental diets and animals (Aghajanzadeh-Golshani et al., 2010; Maheri-Sis et al., 2011b, 2011c). In addition, the use of various equations by different researchers, especially in case of *in vitro* gas production technique, can be resulted in different outputs. Sadeghi & Shawrang (2007) reported that increasing the microwave irradiation time decreased solubility of DM in feedstuffs. The authors supposed that chemical reactions (such as Millard reaction) occurring during heat processing are responsible for the reduction in ruminal degradation. Zhao et al. (2007) stated that microwave irradiation causes biochemical reactions and changes the molecular conformation of starch and protein, texture and physicochemical properties, such as the solubility and gelatin temperature of food products.

Conclusion

The results showed that although microwave irradiation could not improved nutritive value of tomato pomace, but it can be a useful and cost effective method to decrease ruminal gas production (including methane) and removing aflatoxin contamination of this by-product without any negative effects on its nutritional value. It is suggested the use of different microwave power and higher treating time in further studies.

Acknowledgement

This article is adopted from the M.Sc. thesis of Mehdi Eghali-Vaighan in animal science, Islamic Azad University, Shabestar Branch (thesis supervisor: Dr. N. Maheri-Sis). We would like to acknowledge Mr. Jalil Dolgari-Sharaf, Mr. Kiyani Nahand and Mr. Ali Noshadi for their assistance. Authors also thanks to the Animal Research Centre of Islamic Azad University, Shabestar Branch, Iran, and Animal Science Research Institute, Karaj, Iran.

References

Abdollahzadeh F, Pirmohammadi R, Fatehi F, Bernousi I. 2010. Effect of feeding ensiled mixed tomato and apple pomace on

- performance of Holstein dairy cows. Slovak J. Anim. Sci., 43(1): 31-35.
- Aghajanzadeh-Golshani A, Maheri-Sis N, Mirzaei-Aghsaghali A, Baradaran-Hasanzadeh AR. 2010. Comparison of nutritional value of tomato pomace and brewers grain for ruminants using *in vitro* gas production technique. Asian J. Anim. Vet. Adv., 5(1): 43-51.
- Ajmal Khan M, Mahr-Un-Nisa and Sarwar M. 2003. Techniques measuring digestibility for the nutritional evaluation of feeds. Int. J. Agric. Biol., 5(1): 91-94.
- Al-Harashsheha M, Al-Muhtasebb AH, Mageec TRA. 2009. Microwave drying kinetics of tomato pomace: Effect of osmotic dehydration. Chem. Eng. Process., 48: 524-531.
- AOAC. 1990. Official Method of Analysis. 15th Edn., Association of Official Analytical Chemists, Washington, DC., USA., pp: 66-88.
- Besharati M, Taghizadeh A, Janmohammadi H, Moghadam GA. 2008. Evaluation of some by-products using *in situ* and *in vitro* gas production techniques. Am. J. Anim. Vet. Sci., 3: 7-12.
- Chaji, M, Mohammadabadi T, Aghaei A. 2011. Fermenting cell walls of processed sugarcane pith by ruminal bacteria, protozoa and fungi. Int. J. Agric. Biol., 13: 283-286.
- Chavan RS, Chavan SR. 2010. Microwave baking in food industry: a review. Int. J. Dairy Sci., 5: 113-127.
- Chen, Y, Zhao Y, Zhu-Yan F, Zhao-Wei M, Fa-Chen Q, Anwaier A, Bin Y, Reyangu A, Xiao-Li X, Aniwaer A. 2011. Chemical composition and *in vitro* ruminal fermentation characteristics of tetraploid black locust (*Robinia pseudoacacia* L.). Asian J. Anim. Vet. Adv., 6: 706-714.
- Chumpawadee S. 2009. Degradation characteristics of tomato pomace, soybean hull and peanut pod in the rumen using nylon bag technique. Pak. J. Nutr., 8(11): 1717-1721.
- Chumpawadee S, Anut C, Piyanate C. 2007. Chemical compositions and nutritional evaluation of energy feeds for ruminant using *in vitro* gas production technique. Pak. J. Nutr., 6: 607-612.
- Ebrahimi SR, Nikkhah A, Sadeghi AA. 2010. Changes in nutritive value and digestion kinetics of canola seed due to microwave irradiation. Asian-Austral. J. Anim. Sci., 23(3): 347-354.
- Eskicioglu C, Terzian N, Kennedy KJ, Droste RL, Hamoda M. 2007. Athermal microwave effects for enhancing digestibility of waste activated sludge. Water Res., 41: 2457-2466.
- Farag RS, Rashed MM, Abo-Hgger AAA. 1996. Aflatoxin destruction by microwave heating. Int. J. Food Sci. Nutr., 47: 197-208.
- Karn JF. 1991. Chemical composition of forage and feces as affected by microwave oven drying. J. Range Manage., 44: 512-515.
- Kilic U, Garipoglu AV. 2009. *In situ* rumen degradability, *in vitro* digestibility and *in vitro* gas production of full fat canola seeds. Asian J. Anim. Vet. Adv., 4: 200-208.
- Maheri-Sis N, Mirzaei-Aghsaghali A, Mirza-Aghazadeh A, Safaei AR. 2007. Determination of digestibility and nutritive value of Iranian alfalfa varieties using *in vivo* technique in sheep. Asian J. Anim. Vet. Adv., 2(4): 239-243.
- Maheri-Sis, N, Chamani M, Sadeghi AA, Mirza-Aghazadeh A, Aghajanzadeh-Golshani A. 2008. Nutritional evaluation of kabuli and desi type chickpeas (*Cicer arietinum* L.) for ruminants using *in vitro* gas production technique. Afr. J. Biotechnol., 7: 2946-2951.

RESEARCH ARTICLE

- Maheri-Sis N, Baradaran-Hasanzadeh AR, Salamatdoust R, Khosravifar O, Agajanzadeh-Golshani A, Dolgari-Sharaf J. 2011a. Effect of microwave irradiation on nutritive value of sunflower meal for ruminants using *in vitro* gas production technique. *J. Anim. Plant Sci.*, 21(2): 126-131.
- Maheri-Sis N, Eghbali-Vaighan M, Mirza-Aghazadeh A, Ahmadzadeh AR, Aghajanzadeh-Golshani A, Mirzaei-Aghsaghali A, Shaddel-Telli AA. 2011b. Effects of microwave irradiation on ruminal dry matter degradation of tomato pomace. *Curr. Res. J. Biol. Sci.*, 3(3): 268-272.
- Maheri-Sis N, Eghbali-Vaighan M, Mirza-Aghazadeh A, Shaddel-Telli AA, Mirzaei-Aghsaghali A, Aghajanzadeh-Golshani A. 2011c. Effects of microwave irradiation on ruminal protein degradation of tomato pomace. *Int. J. Anim. Vet. Adv.*, 3(3): 177-181.
- Menke KH, Raab L, Salewski A, Steingass H, Fritz D, Schneider W. 1979. The estimation of digestibility and metabolisable energy content of ruminant feedstuffs from the gas production when they incubated with rumen liquor *in vitro*. *J. Agric. Sci.*, 93: 217-222.
- Menke KH, Steingass H. 1988. Estimation of energetic feed value obtained from chemical analysis and *in vitro* gas production using rumen fluid. *Anim. Res. Dev.*, 28: 7-55.
- Mirzaei-Aghsaghali A, Maheri-Sis N. 2008. Nutritive value of some agro-industrial by-products for ruminants: A Review. *World J. Zool.*, 3(2): 40-46.
- Mirzaei-Aghsaghali A, Maheri-Sis N, Mirza-Aghazadeh A, Safaei AR, Houshang AF, Aghajanzadeh-Golshani A. 2008. Use of nylon bag technique to determine nutritive value and degradation kinetics of Iranian alfalfa varieties. *Asian J. Anim. Vet. Adv.*, 3 (4): 214-221.
- Mirzaei-Aghsaghali A, Maheri-Sis N, Mansouri H, Razeghi ME, Safaei AR, Aghajanzadeh-Golshani A, Alipor K. 2011. Estimation of the nutritive value of tomato pomace for ruminant using *in vitro* gas production technique. *Afr. J. Biotechnol.*, 10(33): 6251-6256.
- Ørskov ER, McDonald P. 1979. The estimation of protein degradability in the rumen from incubation measurements weighed according to rate of passage. *J. Agric. Sci.*, 92: 499-503.
- Pelletier S, Tremblay GF, Bertrand A, Bélanger G, Castonguay Y, Michaud R. 2010. Drying procedures affect non-structural carbohydrates and other nutritive value attributes in forage samples. *Anim. Feed Sci. Technol.*, 157: 139-150.
- S.A.S. 1991. SAS for windows, Version 6.0, Ed. SAS Institute Inc., Cary, NC.
- Sadeghi AA, Shawrang P. 2007. Effects of microwave irradiation on ruminal protein degradation and intestinal digestibility of cottonseed meal. *Livest. Sci.*, 106(2-3): 176-181.
- Sadeghi AA, Shawrang P. 2008. Effect of microwave irradiation on ruminal dry matter, protein and starch degradation characteristics of barley grain. *Anim. Feed Sci. Technol.*, 141: 184-194.
- Sindhu AA, Khan MA, Mahr-Un-Nisa, Sarwar M. 2002. Agro-industrial by-products as a potential source of livestock feed. *Int. J. Agric. Biol.*, 4(2): 307-310.
- Sommart K, Parker DS, Rowlinson P, Wanapat M. 2000. Fermentation characteristics and microbial protein synthesis in an *in vitro* system using cassava, rice straw and dried ruzigrass as substrates. *Asian-Austral. J. Anim. Sci.*, 13: 1084-1093.
- Steel RGD, Torrie JH. 1980. Principles and procedures of statistics: a biometrical approach. 2nd Edn., McGraw Hill, New York, USA.
- Taher-Maddah M, Maheri-Sis N, Salamatdoustnobar R, Ahmadzadeh A. 2012. Estimating fermentation characteristics and nutritive value of ensiled and dried pomegranate seeds for ruminants using *in vitro* gas production technique. *Open Vet. J.*, 2: 40-45.
- Tatke P, Jaiswal Y. 2011. An overview of microwave assisted extraction and its applications in herbal drug research. *Res. J. Med. Plant.*, 5: 21-31.
- Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary neutral detergent fiber and non starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74: 3583-3597.
- Zhao S, Xiong S, Qiu C, Xu Y. 2007. Effect of microwaves on rice quality. *J. Stored Products Res.*, 43(4): 496-502.