

RESEARCH ARTICLE

Georgi G. Markov¹
Nuri Yiğit²
Ercüment Çolak²
Maria A. Kocheva¹
Milena K. Gospodinova¹
Hristo Dimitrov³

Sibling vole species (*Microtus*) in the fragmented landscape of south-eastern part of Thrace, Balkan Peninsula: species presence, habitat selection and craniometry

Authors' addresses:

¹ Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 1000 Sofia, Bulgaria

² Biology Department, Faculty of Science, Ankara University, 06100 Beşevler, Ankara, Turkey

³ Department of Zoology, Faculty of Biology, Plovdiv University, 4000 Plovdiv, Bulgaria

Correspondence:

Georgi G. Markov
Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences; 1 Tzar Osvoboditel Blvd., 1000 Sofia, Bulgaria
e-mail: georgimar@gmail.com

Article info:

Received: 29 July 2013

In revised form: 2 November 2013

Accepted: 11 November 2013

ABSTRACT

The presence of sibling vole species (*Microtus*) in the most common forests and open landscapes in south-eastern part of Thrace (Balkan Peninsula) of contemporary territories of Bulgaria and Turkey was studied. Only the southern vole (*Microtus levis*) was found in the investigated region. In his northern part, the Southern vole is associated with semi dry, upland habitats such as deserted and overgrown with wild vegetation vines near sparse forests and large agricultural fields under autumn crops with adjacent non-arable lands with shrub vegetation. In the southern part, it is presented in open landscape related to water areas (meadows near rivers and wetlands). On the basis of trapping the Southern vole is a common species in the north part of Strandzha region. The established craniological characteristics of the Southern vole from investigated region, which can be considered as a zoogeographical crossroads, with a late Pleistocene connection between the Balkan Peninsula and the mammalian fauna of Anatolian peninsula, enriched the knowledge about its craniological variation in Europe.

Key words: *Microtus levis*, southern vole, sibling vole species, south-eastern Thrace, biotopes adherence

Introduction

Population taxonomy of common vole started to change considerably when it was found that the group of the common vole (*Microtus arvalis* sensu lato) in Europe comprises two sibling species: the Common vole (*Microtus arvalis* Pallas, 1778, sensu stricto) with 46 chromosomes and the Southern vole (*Microtus levis* = *rossiaemeridionalis* Ognev, 1924) with 54 chromosomes (Meyer et al., 1969). Examination of karyotypes of numerous common vole populations in Europe, aiming to verify to which one of the sibling species do they belong, revealed widespread sympatric distribution of these two species in their European range (Zima & Král, 1984).

The lack of understanding of the systematic status of many sibling voles' populations in the presumable areas of their sympatric distribution was determined by the fact that

the main diagnostic features of these morphologically similar species, such as karyotype (Meyer et al., 1969) and specific electrophoretic mobility of haemoglobin fractions (Malygin, 1983) were obtained by laborious analyses. The development of craniometrical keys (Markov & Kocheva, 2007, Markov et al., 2009) for species classification of the representatives of *Microtus arvalis* sensu lato, together with the availability of massive craniological material allowed outlining their distribution in many regions in south-east Europe and on the Asiatic part of Turkey.

When it was applied for species craniometric determination of the representatives of *M. arvalis* (sensu lato) on the territory of Bulgaria, it was found that *M. arvalis*, sensu stricto and *M. levis* inhabit together in twenty-three of all studied twenty-five biotopes. Their co-existence is established in North-eastern Bulgaria, Stara Planina foothills, highland fields of Stara Planina, Sredna Gora Mountain and

RESEARCH ARTICLE

Western Bulgaria, sub-Balkan fields and the Black sea coast (Markov & Kocheva, 2008).

At the same time, the distribution and species identity of sibling voles in many other territories of South-eastern Balkan Peninsula, one of which is South-eastern Thrace, have not been clarified and species craniometry was not described. Heretofore no information on species identification of the sibling voles, their habitat distribution and skull shape variation in South-eastern Thrace has ever been reported. Our comprehension of the population taxonomy of the sibling voles (*Microtus*) in Strandzha Mountain region, situated in South-eastern Thrace on the territory of Bulgaria and Turkey, is still insufficient. They are thus far associated with the classic idea about the distribution of the Common vole *M. arvalis*; namely, that only *M. arvalis* could be found there (Niethammer & Krapp, 1978). Detailed population investigations of species identity and craniometry of the sibling voles in South-eastern Thrace on the territory of Turkey (Yiğit *et al.*, 2003) also need filling out.

Thus, the aim of the present work was: (i) to carry out craniometric species identification of the sibling voles' species (*Microtus*) in the south-eastern part of Thrace, (ii) to assess their biotopic adherence particularly in Strandzha Mountain region; and (iii) to characterize craniometrically the identified species.

Materials and Methods

Study area

Animals were collected during summer season (June and July) during the period 2006-2009 in the South-eastern Thrace on the contemporary territories of Bulgaria and Turkey. This part of Balkan Peninsula was connected to Anatolian peninsula by a land bridge during the late Pleistocene (Lang, 1994) and can be considered as a zoogeographical crossroads, with a late Pleistocene connection to the particularly rich mammalian fauna of Anatolian peninsula (Cheylan, 1991). Its mammalian crossroad status is also indicated by the patterns of the present geographical distribution of several terrestrial mammalian species (Niethammer & Krapp, 1978, 1982; Stubbe & Krapp, 1993a, 1993b).

Detailed study of the biotopic adherence of sibling voles (*Microtus*) and their relative abundance in the fragmented landscape of south-eastern part of this region is performed in Strandzha Mountain region. It is situated in south-eastern part of Balkan Peninsula and covers much of South-eastern

Thrace in Bulgaria and the European part of Turkey and belongs to European Deciduous Forest biome (Tahtadzhyan, 1978). The specific geological, climatic and bio-geographical characteristics of this region have determined the formation of natural ecosystems with great biodiversity of small mammals (Paspalev & Markov, 1961; Peshev & Angelova, 1962; Popov, 1993; Markov *et al.*, 2000; Chassovnikarova *et al.*, 2005; Markov *et al.*, 2008). Nowadays, as a result of development of agriculture and grazing, many semi-natural habitats and agricultural areas have been formed (Strandzha Nature Park site), which can be grouped into Woodlands and Open landscapes by their landscape type.

The set of the investigated habitats covering typical and most common plant associations in this region (Bondev, 1991; Gruev & Kuzmanov, 1994) included much of habitats traditionally studied in biodiversity estimation of small mammals in northern Strandzha region (Markov *et al.*, 2000; Chassovnikarova *et al.*, 2005; Markov *et al.*, 2008) and South-eastern Thrace in the European part of Turkey (Yiğit *et al.*, 2003) in both main types landscapes:

(I) Woodlands, represented by:

- (i) Riparian woodlands - periodically flooded riparian mixed deciduous forest (Locations: 11 – Veleka River outfall; 5 – Fakiiska River at Fakiya village);
- (ii) Deciduous woodlands of Colchisian-Mediterranean type (Locations: 10 – Kosti village, 3 – Pismenovo village);
- (iii) Mixed forests, with artificial coniferous plantations (Locations: 6 A – Tsarevo town, 12 – Malko Tarnovo town, 13 – Demirköy village);

(II) Open landscapes, represented by:

- (iv) Meadows near agricultural areas and rivers, covered with steppe-like grass communities, semi-shrub and shrub vegetation (Locations: 6 B – Tsarevo town, 16 – Sarhdere village, 17 – Havsa town; 18 – Tekirdag town);
- (v) Open wetlands (Locations: 9 A – Gramatikovo village; 18 – Tekirdag town);
- (vi) Agricultural lands under alfalfa, autumn cereals or vines (Locations: 8 – Gorno Yabalkovo village, 2 – Iasna Poliana village, 9B – Gramatikovo village; 7 – Ahtopol town; 4 – Novo Panicharevo village);
- (vii) Glades in deciduous and mixed forests (Locations: 6C – Tsarevo town; 15 – Kirkirarli town; 14 – Dereköy village);
- (viii) Pastures with sparse vegetation on drained soil (Locations: 1 – Velika village) (Figure 1).

RESEARCH ARTICLE




Figure 1. Geographic origin of the studied samples from south-eastern part of Thrace. The locations are marked by the names of the nearest settlement: 1- Velika village; 2- Iasna Poliana village; 3 - Pismenovo village; 4 - Novo Panicharevo village; 5 - Fakiya village; 6 A, B, C- Tsarevo town; 7 - Ahtopol town, 8 - Gorno Yabalkovo village; 9 A, B - Gramatikovo village; 10 - Kostı village; 11 - Veleka River outfall; 12 - Malko Tarnovo town; 13 - Demirköy village; 14 - Dereköy village; 15 - Kirklarlı town; 16 - Sarhdere village; 17 - Havsa town; 18 - Tekirdağ town.

The studied complex of habitats covers a significant part of heterogeneous natural-geographic pattern of the South-eastern Thrace and comprises most of the biotopes (both natural and man-made) preferred by the studied sibling vole species. The Southern vole prefers biotopes with tree stands and vegetable gardens, undergrowth of large-stalk grass, etc. (Malygin, 1970, 1983; Dobrokhotov *et al.*, 1985; Tikhonov *et al.*, 1998), while the Common vole typically inhabits coenoses of meadow type and agricultural lands (Dobrokhotov *et al.*, 1985; Teslenko & Zagorodnyuk, 1986; Tikhonov *et al.*, 1992; Karaseva *et al.*, 1994), weakly wooded forest biotopes along the river valleys, ravines, forest belts and avoids territories under intensive human impact and transformation (Tikhonov *et al.*, 1992; Tikhonov *et al.*, 1998; Tikhonov & Tikhonova, 1997).

Collecting and age determination

In each of the studied biotopes the presence of small mammals was surveyed by analysing the species composition of the catch carried out for 2 days by three trap lines each consisting of 25 traps, located 25 meters from each other. The age of the specimens belonging to *M. arvalis* *sensu lato* was determined by craniological criteria (Bashenina, 1953).

Taxonomic classification

All studied adult sibling voles were classified to one of the two sibling species *M. arvalis* or *M. levis* using discriminating craniological function with high classification ability, consisting of four measurements, which did not show sexual dimorphism in the studied species: Alveolar length of upper molars, Zygomatic width, Cranial height between

RESEARCH ARTICLE

bullae ossae and Length of incisive foramen (Markov & Kocheva, 2007). In addition, the method of specific electrophoretic mobility of haemoglobin fractions proposed by Malygin (1983) was applied to verify the species of sibling voles from meadows near agricultural areas and rivers, covered with steppe-like grass communities, semi-shrub and shrub vegetation (Location: Tsarevo town) and agricultural lands under autumn cereals (Location: Ahtopol town); the species of the sibling voles caught in the same biotope (Havsya town; Sarhdere village) or in the wetland biotope (Tekirdag) in Turkish Thrace was verified by karyotyping.

Determining the relative numbers

The relative numbers of the sibling voles in the northern region of south-eastern part of Balkan Peninsula was evaluated in two types of biotopes by the method of trap-lines (Novikov 1953). The first one near the village of Novo Panicharevo village represented neglected and overgrown with wild vegetation vines with sparse deciduous forest nearby and the second near the village of Iasna poliana was represented by agricultural lands under autumn crops. In these two biotopes pitfall traps were used, put in places with clear presence of small mammals – burrows, forming colonies. The presence of small mammals was surveyed by 3 trap lines with 25 traps each for 3 days in the vineyards and by 3 trap lines with 25 traps each for 2 days in the agricultural lands under autumn crops.

The relative number of captured small mammals was determined as number of animals caught per 100 Pitfall days and their dominant presence in the studied area was estimated quantitatively by Kouzyakin's dominance gradation scale (Kouzyakin, 1962) as follows: mass species, if its relative presence is $\geq 30\%$ of the total catch; numerous species, if its relative presence is from 10.0% to 29.9% of the total catch; common species, if its relative presence is from 1.0% to 9.9% of the total catch; rare species, if its relative presence is from 0.2% to 0.9% of the total catch.

Craniometric description

The specimens from Bulgaria and Turkey classified to one of the two species *M. arvalis* or *M. levis* were merged into species' operational taxonomic units for further craniometrical description. Craniometrical analyses were conducted on 42 adult specimens, with established taxonomic affiliation to Southern vole (*M. levis*) from the whole investigated territory of south-eastern part of Thrace. It includes 21 specimens established in the northern part of the

study region on the territory of Bulgaria (Locations: Tsarevo town; Iasna poliana village; Ahtopol town; Novo Panicharevo) and 21 specimens established in its southern part on the territory of Turkey (Locations: Havsya town; Sarhdere village; Tekirdag town).

The set of twenty-six cranial characters (Figure 2), measured with a digital caliper with 0.1 mm accuracy, previously used in obtaining of the craniometric population characteristics of both *Microtus* sibling species in Europe (Markov & Kocheva, 2007) was applied to obtain population craniometrical characteristics and to evaluate species morphometric variation of the studied sibling voles from south-eastern part of Thrace situated on the territory of the Bulgaria and Turkey.

Statistical analysis

All craniometric data were subject to standard statistic univariate analysis. Data were tested for normality using Kolmogorov-Smirnov D-statistics, and for homogeneity of variances using Levene's test. Although previous investigations didn't find any sexual dimorphism in studied characters in the European (Markov & Kocheva, 2007) and Asiatic (Markov *et al.*, 2009) range of the sibling voles, the presence of cranial sexual dimorphism in the studied metric characters was tested by Student's t-test. The probability level for decision about statistical significance was $\alpha = 0.05$. For the overall survey of the skull parameters, the descriptive statistics were counted – the mean (\bar{X}), the standard deviation (SD) and standard error of mean (SE), and the coefficient of variation (CV). The $\pm 95\%$ confidence interval of the mean value of each one variable was also determined. All statistical analyses were carried out using STATISTICA (Version 8.0).

Results

The sibling voles were found only in open landscapes, represented by:

- i. meadows near agricultural areas and rivers, covered with steppe-like grass communities, semi-shrub and shrub vegetation (Locations: Tsarevo town; Havsya town; Sarhdere village);
- ii. open wetlands (Locations: Tekirdag town);
- iii. agricultural lands under alfalfa, autumn cereals or vines (Locations: Iasna Poliana village; Ahtopol town; Novo Panicharevo village).

RESEARCH ARTICLE




Figure 2. Skull and mandible measurements: V1 - Condylbasal length I; V2 - Condylbasal length II; V3 - Basal length; V4 - Condylbasilar length; V5 - Length of incisive foramen; V6 - Total skull length; V7 - Occipito-maxillar length; V8 -Rostrum width; V9 - Zygomatic width; V10 - Occipital width; V11 - Mastoid width; V12 - Interorbital width; V13 - Length of upper incisors; V14 - Thickness of upper incisors; V15 - Rostrum height; V16 - Length of upper diastema; V17 - Height of zygomatic arch; V18 - Alveolar length of upper molars; V19 - Interparietal-foramen magnum height; V20 - Cranial height between bullae ossae; V21 - Cranial height from mastoids V22 - Cranial height from bullae osseae; V23 - Mandible length; V24 - Alveolar length of lower molars; V25 - Articular height; V26 - Mandible height taken at M.

All caught adult sibling voles (*M. arvalis* sensu lato) were craniometrical classified to Southern vole (*M. levis*) (Figure 3). This species identity was confirmed for some individuals by a study of specific electrophoretic mobility of their

haemoglobin fractions (Location: Tsarevo town; Achtopol town) (Markov & Kocheva, 2008); either by establishing their karyotype (Location: Havsa town; Sarhdere village; Tekirdag town) (Yiğit *et al.*, 2003).

RESEARCH ARTICLE




Figure 3. Classification results for taxonomic identification from craniometrical determination of the investigated common vole (*Microtus arvalis*, *sensu lato*) of south-eastern part of Thrace. The values of group centroids are 0.803 for *M. arvalis* and -2.71 for *M. levis*.

The evaluated relative numbers of the sibling voles in the two types of biotopes in the northern region of south-eastern part of Balkan Peninsula showed that:

- i. In biotope represented by neglected and overgrown with wild vegetation vines with sparse deciduous forest nearby (near the village of Novo Panicharevo), except the sibling voles, classified craniometrically as *M. levis*, the yellow-necked field mouse (*Apodemus flavicollis*) and bicolored white-toothed shrews (*Crocidura leucodon*) were also found. The relative numbers of the southern vole in this biotope was 2.65 individuals per 100 trap-days, while the relative numbers of the yellow-necked field mouse was 5.31 individuals per 100 trap-days, and the bicolored white-toothed shrew was presented only by 0.44 individuals per 100 trap-days.
- ii. In the other biotope, represented by agricultural lands under autumn crops (Iasna poliana village) all captured voles were also classified as *M. levis*. In this biotope, the relative numbers of the southern vole was 4.0 individuals per 100 trap-days. The same relative numbers of 4.0 individuals per 100 trap-days demonstrated the second species, which was found to inhabit this biotope - the yellow-necked field mouse.

The differences between each cranial measurements of adult male and female specimens of Southern vole were not significant ($p > 0.05$, Student's *t*-test) in all studied variables and the obtained varied from $t = 0.20$ to $t = 1.68$. The lack of

statistical evidence of differences in the average values of the studied parameters in both sexes of Southern vole allow them to be pooled together for further craniometrical analyses. Based on the classified specimens, a population craniometrical characteristic of the southern vole in south east part of Balkan Peninsula (south-eastern Thrace region) was made (Table 1). All studied craniometrical characters were identified with similar relatively low standard error of the mean and they all showed similar absolute variation. The confidence limits for the means of all investigated craniometrical parameters limited too narrow range of values around the means, where the population means are expected to be located.

Phenotype craniological variation of the southern vole from the south-eastern part of Thrace assessed by comparing the relative variation (CV) of the studied craniometrical characters was within the range from 3.5% to 10.0% (Table 1), which is typical for the most rodents.

Discussion

Taxonomic determinations of the established representatives of *M. arvalis sensu lato* evidently revealed the absence of Common vole (*M. arvalis sensu stricto*) and confirmed that the Southern vole is presented in the fragmented landscape of south-eastern part of Thrace.

In northern Strandzha mountain region the Southern vole is associated with semi dry, upland habitats such as deserted and overgrown with wild vegetation vines near sparse forests and large agricultural fields under autumn crops with adjacent non-arable lands with shrub vegetation. These biotopes correspond to the ones preferred by the species in other parts of its European range. In this region the sibling species *M. arvalis sensu stricto* was not established by trapping, in the biotopes considered as suitable for its development, such as coenoses of meadow type, agricultural lands under alfalfa or natural biotopes with sparse shrub vegetation along the river valleys. Although the relative numbers of the Southern vole remains near the lower average limit of the grade estimation for presence of common species in particular biotope, the species still could be considered as common in the biotopes, where it was found.

At the same time in the south part of Thrace, Southern vole is presented in open landscapes related to water areas (meadows near rivers and wetlands). This fact suggested an ubiquitous colonization of south-eastern part of Thrace by the Southern vole.

RESEARCH ARTICLE

Table 1. Descriptive statistic of investigated craniometrical variables of Southern vole (*Microtus levis*) from south-eastern part of Thrace. Skull and mandible measurements are described on Figure 2.

	Valid N	Mean \bar{X}	Standard error (SE)	Standard deviation (SD)	+95% Confid. interval	-95% Confid. interval	Coef. of Var. (CV)
V1	38	24.59	0.213	1.311	24.16	25.02	5.3
V2	38	25.10	0.202	1.246	24.69	25.51	4.9
V3	35	22.43	0.210	1.241	22.00	22.86	5.5
V4	38	23.93	0.216	1.329	23.49	24.36	5.5
V5	42	4.46	0.046	0.301	4.37	4.56	6.7
V6	41	25.10	0.193	1.233	24.71	25.49	4.9
V7	41	19.68	0.140	0.895	19.40	19.97	4.5
V8	42	4.20	0.049	0.316	4.10	4.30	7.5
V9	42	14.24	0.138	0.891	13.96	14.52	6.2
V10	42	11.49	0.089	0.579	11.31	11.67	5.0
V11	42	10.59	0.085	0.551	10.41	10.76	5.2
V12	42	3.68	0.024	0.156	3.64	3.73	4.2
V13	41	4.49	0.053	0.341	4.39	4.60	7.5
V14	42	1.48	0.020	0.127	1.44	1.52	8.5
V15	42	4.14	0.045	0.289	4.05	4.23	6.9
V16	42	7.46	0.066	0.425	7.32	7.59	5.7
V17	42	1.52	0.023	0.149	1.48	1.57	9.8
V18	42	6.29	0.034	0.223	6.22	6.36	3.5
V19	40	3.30	0.050	0.317	3.20	3.40	9.6
V20	42	7.40	0.068	0.438	7.27	7.54	5.9
V21	42	9.27	0.062	0.400	9.15	9.40	4.3
V22	42	7.54	0.050	0.321	7.44	7.64	4.2
V23	40	15.31	0.112	0.707	15.08	15.53	4.6
V24	42	5.97	0.049	0.320	5.87	6.07	5.3
V25	19	7.98	0.109	0.474	7.75	8.21	5.9
V26	41	4.75	0.036	0.231	4.68	4.83	4.8

The established population craniometrical characters of *M. levis* from Southeastern Balkan Peninsula, a region considered as a zoogeographical crossroads and with key geographic position in the species range, could be used as baseline for estimation of bio-morphological diversity of this species in its Eurasian range. The evaluation of craniological variation and local craniometrical characteristics of *M. levis* under the specific ecological conditions of the south-eastern part of Thrace, particularly in Strandzha Mountain region, affords an opportunity for comparative analysis and revealing of craniological population diversity of this species in main natural and agricultural ecosystems situated on the territory of the Bulgaria and Turkey.

The unambiguously determined relative and absolute variation of craniometrical characters of *M. levis* from this region expands significantly the potential in solving many problems related to assessment of its craniological populations variation. Moreover, such craniological

characteristics could be expanded when the species is used as bio-indicator as an integral part of bio-monitoring system of rating environment quality (Peev & Gerasimov, 1990) in south-eastern part of Thrace, both in the two countries and in other specific parts of the species range.

Acknowledgement

This research was partially supported by the Scientific and Technical Research Council of Turkey (TUBITAK) within 2221-Visiting Scientist Fellowship Program.

References

- Bashenina NV. 1953. On age determination of common vole (*Microtus arvalis* Pall.). *Zool. Zh.*, 32 (4): 730-743 (In Russian).
 Bondev I. 1991. Vegetation in Bulgaria. – St. Kl. Ohridski University Press, Sofia, 183 p. (In Bulgarian).

RESEARCH ARTICLE

- Chassovnikarova T, Dimitrov H, Markov G, Mitev D. 2005. Distribution and habitat association of field mouse (*Apodemus agrarius* Pallas, 1771) in National Park "Strandja". In: Proc. of the Balkan Scientific conference of biology in Plovdiv (Bulgaria) 19-21 May 2005 (Eds. Gruev B, Nikolova M, Donev A), 428-436.
- Cheylan G. 1991. Patterns of Pleistocene turnover, current distribution and speciation among Mediterranean mammals. – In: Groves RH & FDi Castri (eds.), *Biogeography of Mediterranean Invasions*, Cambridge University Press, Cambridge, 227-262.
- Dobrokhotov BP, Baranovskii PM, Demidova TN. 1985. Peculiarities of station distribution of the common vole sibling species *Microtus arvalis* and *Microtus rossiaemerdionalis* (Rodentia, Cricetidae). *Zool. Zh.*, 64(2): 269-275 (in Russian, English summary).
- Gruev B, Kuzmanov B. 1994. *General biogeography*. – St. Kliment Ohridski University Press, Sofia, 498 p. (in Bulgarian).
- Karaseva EV, Stepanova NV, Telicyna AYU, Merzlikin IR, Posel'skaya OI. 1994. Ecological differences of two sibling species – the common and East-European voles – In: *Synanthropy of rodents*, IEMEZH RAN, Moscow, p. 60-74. (In Russian).
- Kuzyakin AP. 1962. *Zoogeography USSR*. – MOPI. C., IX, I: 3-157
- Lang G. 1994. Quartäre Vegetationsgeschichte Europas. *Methoden und Ergebnisse*. – G. Fischer, Jena: 1-462.
- Malygin VM. 1970. Systematic of super-species of common vole. *Vestnik MGU, Ser. Biol.*, 5: 89-91. (In Russian)
- Malygin VM. 1983. *Systematics of the Common Vole*. Moscow, 207 pp. (in Russian)
- Markov G. 1959. *Mammals in Bulgaria*. – Science and Art, Sofia, 155 p. (in Bulgarian).
- Markov G, Mitev D, Chassovnikarova T, Dimitrov H. 2008. Habitat heterogeneity and small mammal species richness in the Strandja nature park (SE Bulgaria): Implications for conservation. *Mammalian Biology*, 73: 25.
- Markov G, Atanasov N, Chassovnikarova Ts. 2000. Ecological faunistic review and population biological analysis of Micromammalia in Strandja national Park. – In: *Proceedings of International Conference "75 Years of Higher Forestry Education in Bulgaria"*, Sofia. 67-74. (in Bulgarian)
- Markov G, Yigit N, Kocheva M, Colak E. 2009. A refine method for craniometrical identification of the sibling vole species *Microtus arvalis* and *Microtus rossiameridionalis* in Europe and the Asiatic part of Turkey, *N-W J. Zool.*, 5(1): 1-7.
- Markov G, Kocheva M. 2007. Craniometric characteristics and differentiation of the sibling vole species *Microtus arvalis* and *Microtus rossiaemerdionalis*. *Acta Zool. Bulg.*, 59(2): 145-152.
- Markov G, Kocheva M. 2008. Sympatric distribution of the sibling vole species *Microtus arvalis* and *Microtus levis* (Rodentia: Cricetidae) in Bulgaria. *Acta Zool. Bulg.*, 60(1): 107-111.
- Meyer MN, Orlov VN, Skholl' ED. 1969. New rodent species (Rodentia, Mammalia) found on the grounds of karyological, physiological and cytophysiological analyses. *Dokl. AN USSR*, 188(6): 1411-1414. (In Russian)
- Niethammer J, Krapp F. (eds) 1978. *Handbuch der Säugetiere Europas*. Vol. I/Nagetiere 1. – Akademische Verlagsgesellschaft Wiesbaden: 1-476.
- Niethammer J, Krapp F. (eds) 1982. *Handbuch der Säugetiere Europas*. Vol. 2/Nagetiere II. – Akademische Verlagsgesellschaft Wiesbaden: 1-649.
- Novikov GA. 1953. *Field studies on the ecology of terrestrial vertebrates*. – Soviet science, 501 pp. (In Russian).
- Paspalev G, Markov G. 1961. Ecological faunistic investigations on Mammals in Strandzha Mountain. – In: *Natural diseases sources in Strandzha Mountain Region (Proc.)*. BAS, 3-20. (In Bulgarian).
- Peshev Ts, Angelova V. 1962/1963. Insectivora and Rodents in Southeastern outskirts of Bulgaria. *Annals of Sofia University, LVII (1), Biology (Zoology)*, 69-75. (In Bulgarian).
- Popov V. 1993. Small mammals (Insectivora, Lagomorpha, Rodentia). – In: *National strategy for protection of biological diversity*, I: 615-630. (In Bulgarian).
- Peev D, Gerasimov S. (eds.) 1990. *National program for biomonitoring of Bulgaria*. – Gea Libris, Sofia, pp. 240
- Strandja Nature Park site <http://www.strandja.bg/BG/steady-tourism_bg.html>
- Stubbe M, Krapp F. 1993a. *Handbuch der Säugetiere Europas*. Raubsäuger Teil I. – Aula Verlag, Wiesbaden: 1-526.
- Stubbe M, Krapp F. 1993b. *Handbuch der Säugetiere Europas*. Raubsäuger Teil I. – Aula Verlag, Wiesbaden: 529-1213.
- Tahtadzhyan AL. 1978. *Floristic regions of Earth*. – Nauka, Moscow. (in Russian)
- Teslenko SV, Zagorodnyuk IV. 1986. Sibling species of the *Microtus arvalis* super-species in the Ukraine. *Communication II. Occurrence of Microtus arvalis*. *Vest. Zool.*, 6: 27-31 (in Russian).
- Tikhonov IA, Tikhonova GN. 1997. Diversity and perspectives for survival of the voles of genus *Microtus* on urbanized territories. *Proceedings of the Conference on Dynamics of biodiversity of animals' world*, p. 107-111. (in Russian).
- Tikhonov IA, Tikhonova GN, Karaseva EV. 1992. Small mammals in rural settlements of the Middle Russia. – In: *Synanthropy of rodents and control on their numbers*. Moscow, Russian Academy of Sciences. p. 333-354. (in Russian)
- Tikhonov IA, Tikhonova GN, Polyakova LV. 1998. Sibling species of *Microtus arvalis* and *Microtus rossiaemerdionalis* (Rodentia, Cricetidae) in the northeastern Moscow district. *Zool. Zh.*, 77(1): 95-100. (in Russian)
- Yiğit N, Colak E, Sozen M, Ozkurt Ş. 2003. A study on the geographic distribution along with habitat aspects of rodent species in Turkey. *Bonn. Zool. Beitr.*, 50: 355-368.
- Zima J, Král B. 1984. Karyotypes of European mammals II. *Acta Sci. Nat. Acad. Sci. Bohem., Brno*, 18(8) 1-62.