

RESEARCH ARTICLE

Sonia Moussouni¹
Dimitris P. Makris²

Optimisation of onion peroxidase-catalysed formation of aureusidin using 2',4',6',3,4-pentahydroxy chalcone as substrate

Authors' addresses:

¹ Food Quality & Chemistry of Natural Products, Mediterranean Agronomic Institute of Chania (M.A.I.Ch.), Centre International de Hautes Etudes Agronomiques Méditerranéennes, Chania - 73100, GREECE.

² Department of Food Science & Nutrition, University of the Aegean, Myrina - 81400, Lemnos, GREECE.

Correspondence:

Dimitris P. Makris
Department of Food Science & Nutrition, University of the Aegean, Mitr. Ioakim Street, Myrina - 81400, Lemnos, GREECE
Tel.: +30 22540 83114
e-mail: dmakris@aegean.gr

Article info:

Received: 20 March 2014

Accepted: 9 June 2014

ABSTRACT

Previous investigations demonstrated that crude peroxidase (POD) obtained from onion solid wastes has the ability to catalyse the formation of the aurone aureusidin (ARS), using 2',4',6',3,4-pentahydroxy chalcone (PHC) as substrate, although this reaction under physiological conditions is mediated by a polyphenol oxidase-like enzyme, called aureusidin synthase (AS). In this study, a crude onion POD preparation was used to study the effect of some critical factors affecting the reaction, including reaction time, pH and temperature. The optimal set of conditions was identified by deploying central composite factorial design and response surface methodology. The results obtained showed that the optimum values for pH and temperature were 5 and 20°C, respectively, while time was found to exert a statistically non-significant effect. These values were the same or very close to optimal conditions found for structurally different onion POD substrates. The outcome was discussed with regard to the applicability of the onion POD as a versatile tool of biocatalysis.

Key words: aurones, aureusidin, onion, peroxidase, response surface methodology

Abbreviations:

ARS, aureusidin; AS, aureusidin synthase; PHC, 2',4',6',3,4-pentahydroxy chalcone; POD, peroxidase; PPO, polyphenol oxidase; THC, 2',6',3,4-tetrahydroxy chalcone; BSA - bovine serum albumin.

Introduction

Flavonoids are a widely occurring group of polyphenols, embracing several thousands of structures, with versatile biological utility. Prominent members are flavanols (catechins and condensed tannins), flavonols, anthocyanins (water-soluble red pigments) and flavanones, although there is a bewildering diversity in the relevant amounts and width of occurrence (Haminiuk et al., 2012). Apart from their activities pertaining to some vital functions of plants (Treutter, 2006), flavonoids have attracted substantial interest due to their multiple biological effects and their significant potency as pharmacological agents (Curin & Andriantsitohaina, 2005). In this context, numerous studies

have been carried out, revealing a spectrum of bioactivities and pharmacological properties (Dai & Mumper, 2010).

Aurones represent a minor category of flavonoids, both in terms of occurrence and burden in plant tissues, yet their biological properties might be of significant value, as exemplified by recent reports (Lawrence et al., 2003; Detsi et al., 2009). In this regard, their examination with respect to structure, synthesis and activity merits greater attention. In plant tissues, aurones are biosynthesised by chalcone precursors, with the mediation of an enzyme called aureusidin synthase (AS) (Nakayama et al., 2001; Ono et al., 2006). This peculiar biocatalyst has been characterised as a chalcone-specific polyphenol oxidase (PPO) homologue, while its action upon classic PPO substrates is either particularly low or inexistent (Nakayama et al., 2001). The reaction catalysed by AS has been proposed to involve three steps, only the first one being enzyme-dependent (Figure 1) (Nakayama et al., 2002). In this step, the enzyme-catalysed reaction generates a chalcone with an *o*-quinone form of the

RESEARCH ARTICLE

B-ring [intermediate (a)] from 2',6',3,4-tetrahydroxy chalcone (THC) or PHC. The activated double bond of this intermediate undergoes then nucleophilic attack by its 2'-hydroxyl group to yield the intermediate (b), which in turn undergoes further rearrangement (tautomerism) to produce the aurone.




Figure 1. Proposed mechanism of aurone synthesis from PHC catalysed by aureusidin synthase (AS).

The chemical synthesis of aurones has been based mainly on inorganic catalytic means (Bose et al., 2001; Detsi et al., 2009; Lawrence et al., 2003) but recently, the enzymic synthesis of ARS, using PHC as substrate and a crude peroxidase prepared from onion solid wastes has been reported (Moussouni et al., 2010). In extending that study, the investigation presented herein is a deeper insight into ARS synthesis using a crude onion POD. The optimal set of conditions, regarding reaction time, pH and temperature, was identified by deploying central composite factorial design and response surface methodology.

Materials and Methods

Chemicals

All solvents used for chromatography were HPLC grade. 2',3,4,4',6'-pentahydroxy-chalcone (PHC) was from Extrasynthèse (Geny, France). Quercetin was from Sigma Chemical Co (St. Louis, MO, U.S.A.). Hydrogen peroxide (30%), *N,N*-dimethylformamide (DMF) and trichloroacetic acid (TCA) were from Merck (Germany). For all pH values tested, a phosphate/citrate buffer was used.

Preparation of the onion solid waste homogenate

The protocol previously described was used (Osman et al., 2008). Briefly, onion solid wastes were ground in a domestic blender and an aliquot of 2 g of the ground tissue was suspended in 15 mL buffer solution under stirring. The suspension formed was centrifuged at 3,000×*g* for 20 min and then filtered through paper filter to remove cell debris. The clear supernatant obtained was treated with activated charcoal for decolourisation, and filtered through celite under vacuum. The clear filtrate was used as the crude enzyme source.

Peroxidase activity

A previously described methodology was used (Osman et al. 2008), to determine the POD activity of the crude preparations. An aliquot of 0.1 mL of quercetin (0.1 mM in DMF) solution was mixed with 0.1 mL crude extract and 0.8 mL H₂O₂, and the reaction was followed by measuring the decrease in absorbance at 370 nm. One enzyme unit (U) was defined as ΔA₃₇₀ s⁻¹. Control reactions by omitting H₂O₂ or using heat-inactivated crude extract were also carried out. In assays performed at different temperatures, all constituents of the reaction mixture were pre-incubated in a thermostated water bath. Protein content of the homogenate was determined according to Bradford (1976), using BSA as standard. For all determinations, a computer-controlled HP 8452A diode-array spectrophotometer was used.

Aureusidin synthesis

Stock solutions of PHC (17.3 mM) were prepared in ethanol. PHC solution was diluted 1/10 with DMF prior to reactions. The reaction mixture was composed of 0.8 mL citrate/phosphate buffer (pH 3, 5 or 7) containing 3 mM H₂O₂, 0.1 mL crude enzyme extract (6.5 mU) and 0.1 mL of the diluted PHC solution. The mixture was incubated at various temperatures for predetermined periods, as dictated by the experimental design (Table 1). The reaction was arrested by adding 0.1 mL TCA (10% w/v in ethanol) and centrifuging the mixture at 5,000×*g* for 10 min.

Liquid chromatography-mass spectrometry (LC-MS)

The LC-MS analysis was performed using an LC-DAD-MS system comprising a Finnigan MAT Spectra System P4000 pump, coupled with a UV6000LP diode array detector and a Finnigan AQA mass spectrometer. The separation was performed on a 125×2 mm Superspher 100-4, RP-18 column (Macherey-Nagel, 4 μm particle size) at a flow rate of 0.33 mL min⁻¹, the column being kept at 40°C. The detection was

RESEARCH ARTICLE

monitored at 278 nm. The MS-ESI(+) spectroscopy was performed at a probe temperature of 350°C, probe voltage of 4 kV and at 12 and 40 eV collision energy in the mass analyzer. The mass range was set at 121–787 amu and the scan rate was 0.8 scans s⁻¹. The following gradient programme was used: (A) acetic acid (2.5%) and (B) methanol; 100% A for 5 min, 0% A in 15 min and kept at 0% A for another 5 min. The data were processed using the Xcalibur 1.2 software.

Table 1. Experimental values and coded levels of the independent variables used for the 2³ full-factorial design.

Independent variables	Code units	Coded variable level		
		-1	0	1
T (°C)	X ₁	20	30	40
pH	X ₂	5	6	7
Time (h)	X ₃	1	3	5

Experimental design and statistical analyses

A 2³ full-factorial, central composite experimental design was used to identify the relationship existing between the response function and process variables, as well as to determine those conditions that optimised the POD-catalysed process. The response function considered was the ARS production, estimated by the peak area at 278 nm, in the chromatograms obtained by the LC-MS system. The three independent variables or factors considered were temperature (X₁, varying between 20 and 40°C), pH (X₂, varying between 3 and 7) and time [X₃, varying between 1 and 5 h]. Each variable to be optimised was coded at three levels, -1, 0 and 1 (Table 1). The choice of value ranges for each variable was based on preliminary experimentation and literature (Moussouni et al., 2010).

The three independent variables were coded according to the following equation:

$$x_i = \frac{X_i - X_0}{\Delta X_i}, \quad x_i = 1, 2, 3$$

where x_i and X_i are the dimensionless and the actual value of the independent variable *i*, X₀ the actual value of the independent variable *i* at the central point, and ΔX_i the step change of X_i corresponding to a unit variation of the dimensionless value. Response at each design point was

recorded (Table 2). Data from the central composite experimental design were subjected to regression analysis using least square regression methodology to obtain the parameters of the mathematical models.

The statistical significance of the regression coefficients deriving from the model was assessed by Student's *t*-test. Analysis of variance (ANOVA) was used to evaluate the overall statistical significance of the model. Response surface plots were obtained using the fitted model, by keeping the independent variables simultaneous. All determinations were carried out at least in triplicate and values were averaged and given along the standard deviation (± S. D.). For all statistics, Microsoft Excel™ 2000, SigmaPlot™ 11 and JMP™ 8 were used.

Table 2. Measured and predicted ARS peak area values determined for individual design points.

Design point	Independent variables			Response	
	X ₁	X ₂	X ₃	Peak area (aureusidin)	
				Measured	Predicted
1	-1	-1	-1	277.8	283.1
2	-1	-1	1	324.8	308.7
3	-1	1	-1	0	0
4	-1	1	1	0	0
5	1	-1	-1	136.6	150.3
6	1	-1	1	156.1	180.8
7	1	1	-1	0	14.3
8	1	1	1	37.5	30.3
9	-1	0	0	0	52.9
10	1	0	0	54.9	9.4
11	0	-1	0	292.5	265.0
12	0	1	0	0	34.9
13	0	0	-1	0	0
14	0	0	1	0	14.1
15	0	0	0	49.3	34.5
16	0	0	0	38.4	34.5

Results and Discussion

Tentative identification of the reaction product

In Figure 2 can be seen a chromatographic trace recorded at 278 nm. The compound eluted at 8.28 min could easily be identified as the starting material (PHC), based on the retention time of an original standard. The peak eluted at 7.75 min showed a typical aurone UV-vis spectrum, with λ_{max} at 398 nm and two shoulders at 268 and 322 nm (Figure 3). The mass spectrum revealed a pseudo-molecular ion at *m/z* = 287

RESEARCH ARTICLE

and diagnostic ions at $m/z = 309$ (Na^+ adduct) and $m/z = 595$, which could be ascribed to a Na^+ adduct of aureusidin dimer $[2\text{M}+23]^+$. Thus this peak was assigned to ARS.

Optimisation of reaction conditions

Values of the independent process variables (X_1 , X_2 and X_3) considered, as well as measured and predicted values for the response (aureusidine production as peak area) are analytically given in Table 2. The experimental values of the response were analysed by multiple regression to fit the following second-order polynomial equation:

$$\text{Peak area} = 34.54 - 21.75X_1 - 115.03X_2 + 10.4X_3 + 43.43X_1X_2 + 1.25X_2X_3 - 3.62X_1X_3 - 30.41X_1^2 + 115.39X_2^2 - 30.86X_3^2$$

The quality of fit was ascertained using the regression coefficients (R^2). The experimental data obtained showed a good fit with the equations ($R^2 = 0.95$, $p = 0.0083$). This fact indicated a satisfactory agreement between observed and predicted responses and that the equation found can adequately predict the experimental results.

After removal of the non-significant factors, as revealed by the ANOVA analysis, the theoretical model could be simplified as follows:

$$\text{Peak area} = 34.54 - 115.03X_2 + 43.43X_1X_2 + 115.39X_2^2$$

The utilisation of the predictive model enabled the theoretical calculation of the optimal set of conditions, that were $\text{pH} = 5$, $t = 1$ h, and $T = 20$ °C. Under these conditions, the maximum theoretically calculated peak area was 328.7 ± 83.0 . The trend concerning the co-variation of the two significant variables (T , pH) was recorded in the form of a three-dimensional plot (Figure 4).

The optimum pH value is in accordance with previous findings on onion POD for a variety of substrates, including the flavonol morin (Osman & Makris, 2011), as well as simpler molecules, such as caffeic acid (El Agha et al., 2009) and *p*-coumaric acid (El Agha & Makris, 2012). In addition, ARS production upon AS action on THC as substrate was found to have an optimum pH 5.4 (Sato et al., 2001). Likewise, the optimum temperature is in line with that found for chlorogenic acid (Osman et al., 2012), but also close to the values reported for the flavonols fisetin (Osman & Makris, 2010) and morin (Osman & Makris, 2011) and for caffeic acid (El Agha et al., 2009). Based on this information and the data presented herein, it would appear that the crude onion POD exhibits similar biochemical characteristics with a variety of substrates. However, optimisation using response surface methodology enabled the identification of the ideal

combination of variables that favoured high turnover rates for the specific reaction of PHC oxidation into ARS. The latter remains to be elucidated by kinetic studies.




Figure 2. An HPLC trace recorded at 278 showing both the starting material (PHC) and the product (ARS).




Figure 3. UV-Vis (upper figure) and mass spectrum (lower figure) of ARS.

Peroxidases are considered versatile biocatalytic tools, mediating a range of oxidative reactions and have been used for purposes of bioremediation (Hamid & Rahman, 2009; Barakat et al., 2010), synthesis of chemicals, as well as generation of bio-based substances, including sinapic acid dimers and other oligomers (Liu et al., 2007) and ferulic

RESEARCH ARTICLE

acid/resveratrol heterodimers (Yu et al., 2007). Peroxidases have also been shown to form ferulic and caffeic acid dimers (El Agha et al., 2008a; 2009) and similar products from chlorogenic and *p*-coumaric acids (El Agha & Makris, 2012; Osman et al. 2012).

However, quercetin oxidation by crude onion POD was demonstrated to afford oxidation products not encountered in reactions with peroxidases from other sources (Osman et al., 2008). Further to that, there has been substantial evidence that onion POD can act in the same manner with PPO, producing structurally similar compounds, as exemplified by the case of ARS formation using PHC as substrate. Recent reports also showed that potato peel PPO-catalysed hydrocaffeic acid oxidation (Demian & Makris, 2013) yielded exactly the same products as the onion POD-catalysed reaction (El Agha et al., 2008b). The fact that crude onion POD might use alternative pathways for substrate oxidation merits a profounder examination, as it may lead to novel applications in green synthesis and other biocatalytic processes.




Figure 4. Response surface plot showing the effect of temperature (X_1) and pH (X_2) co-variation on ARS production.

Conclusion

A crude onion POD was used to generate ARS using PHC as substrate. Critical parameters that affect the enzymic reaction, including temperature, pH and time, were assessed simultaneously by deploying central composite factorial design and response surface methodology. The optimum values for ARS production regarding temperature and pH

were 20°C and 5, respectively. Time was found to exert a statistically non-significant effect and thus the shorter interval tested (1 h) was chosen. The findings concerning pH and temperature were consistent with previous reports on other, structurally unrelated substrates, which suggested that the enzyme preparation expresses optimal activity within narrow ranges of conditions. It is proposed that kinetic studies are required to identify dependence of activity on both PHC and H_2O_2 concentrations, as well as to illuminate putative mechanism of action. This is particularly important, given the sound evidence that the oxidation mechanism(s) involved might share common features with the ones proposed for polyphenol oxidases.

References

- Barakat N, Makris DP, Kefalas P, Psillakis E. 2010. Investigation on the removal of olive mill waste water phenolics with the use of a crude peroxidase extract from onion solid by-products. *Environ. Chem. Lett.*, 8: 271-275.
- Bose G, Mondal E, Khana AT, Bordoloi MJ. 2001. An environmentally benign synthesis of aurones and flavones from 2'-acetoxychalcones using *n*-tetrabutylammonium tribromide. *Tetrahydron Lett.*, 42: 8907-8909.
- Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- Curin Y, Andriantsitohaina R. 2005. Polyphenols as potential therapeutic agents against cardiovascular diseases. *Pharmacol. Rep.*, 57(s): 97-107.
- Dai J, Mumper RJ. 2010. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules* 15: 7313-7352.
- Demian FD, Makris DP. 2013. Factorial design optimisation of hydrocaffeic acid removal from an aqueous matrix by the use of a crude potato polyphenol oxidase. *Biocatal. Agric. Biotech.*, 2: 305-310.
- Detsi A, Majdalani M, Kontogiorgis CA, Hadjipavlou-Litina D, Kefalas P. 2009. Natural and synthetic 2'-hydroxy-chalcones and aurones: Synthesis, characterization and evaluation of the antioxidant and soybean lipoxygenase inhibitory activity. *Bioorgan. Med. Chem.*, 17: 8073-8085.
- El Agha A, Makris DP, Kefalas P. 2008a. Peroxidase-active cell free extract from onion solid wastes: biocatalytic properties and putative pathway of ferulic acid oxidation. *J. Biosci. Bioeng.*, 106: 279-285.
- El Agha A, Makris DP, Kefalas P. 2008b. Hydrocaffeic acid oxidation by a peroxidase homogenate from onion solid wastes. *Eur. Food Res. Technol.*, 227: 1379-1386.
- El Agha A, Abbeddou S, Makris DP, Kefalas P. 2009. Biocatalytic properties of a peroxidase-active cell-free extract from onion solid wastes: caffeic acid oxidation. *Biodegradation*, 20: 143-153.

RESEARCH ARTICLE

- El Agha A, Makris DP. 2012. Biocatalytic characteristics, product formation and putative pathway of *p*-coumaric acid oxidation by a crude peroxidase from onion. *Acta Aliment.*, 41: 304-315.
- Hamid M, Rahman K-u. 2009. Potential applications of peroxidases. *Food Chem.*, 115: 1177-1186.
- Haminiuk CWI, Maciel GM, Plata-Oviedo MSV, Peralta RM. 2012. Phenolic compounds in fruits – an overview. *Int. J. Food Sci. Technol.*, 47: 2023-2044.
- Lawrence NJ, Rennison D, McGown AT, Hadfield JA. 2003. The total synthesis of an aurone isolated from *Uvaria hamiltonii*: aurones and flavones as anticancer agents. *Bioorg. Med. Chem.*, 13: 3759-3763.
- Liu H-L, Wan X, Huang X-F, Kong L-Y. 2007. Biotransformation of sinapic acid catalyzed by *Momordica charantia* peroxidase. *J. Agric. Food Chem.*, 55: 1003-1008.
- Moussouni S, Detsi A, Majdalani M, Makris DP, Kefalas P. 2010. Crude peroxidase from onion solid waste as a tool for organic synthesis. Part I: Cyclisation of 2',3,4,4',6'-pentahydroxy-chalcone to aureusidin. *Tetrahedron Lett.*, 51: 4076-4078.
- Nakayama T, Sato T, Fukui Y, Yonekura-Sakakibarab K, Hayashi H, Tanaka Y, Kusumi T, Nishino T. 2001. Specificity analysis and mechanism of aurone synthesis catalyzed by aureusidin synthase, a polyphenol oxidase homolog responsible for flower coloration. *FEBS Lett.*, 499: 107-111.
- Nakayama T. 2002. Enzymology of aurone biosynthesis. *J. Biosci. Bioeng.*, 94: 487-491.
- Ono E, Hatayama M, Isono Y, Sato T, Watanabe R, Yonekura-Sakakibara K, Fukuchi-Mizutani M, Tanaka Y, Kusumi T, Nishino T, Nakayama T. 2006. Localization of a flavonoid biosynthetic polyphenol oxidase in vacuoles. *Plant J.*, 45: 133-143.
- Osman A, Makris DP, Kefalas P. 2008. Investigation on biocatalytic properties of a peroxidase-active homogenate from onion solid wastes: an insight into quercetin oxidation mechanism. *Process Biochem.*, 43: 861-867.
- Osman A, Makris DP. 2010. Comparison of fisetin and quercetin oxidation with a cell-free extract of onion trimmings and peel, plant waste, containing peroxidase enzyme: a further insight into flavonol degradation mechanism. *Inter. J. Food Sci. Technol.*, 45: 2265-2271.
- Osman A, Makris DP. 2011. Oxidation of morin (2',3,4',5,7-pentahydroxyflavone) with a peroxidase homogenate from onion. *Inter. Food Res. J.*, 18: 1039-1043.
- Osman A, El Agha A, Makris DP, Kefalas P. 2012. Chlorogenic acid oxidation by a crude peroxidase preparation: biocatalytic characteristics and oxidation products. *Food Bioproc. Technol.*, 5: 243-251.
- Sato T, Nakayama T, Kikuchi S, Fukui Y, Yonekura-Sakakibara K, Ueda T, Nishino T, Tanaka Y, Kusumi T. 2001. Enzymatic formation of aurones in the extracts of yellow snapdragon flowers. *Plant Sci.*, 160: 229-236.
- Treutter D. 2006. Significance of flavonoids in plant resistance: a review. *Env. Chem. Lett.*, 4: 147-157.
- Yu B-B, Han X-Z, Lou H-X. 2007. Oligomers of resveratrol and ferulic acid prepared by peroxidase-catalyzed oxidation and their protective effects on cardiac injury. *J. Agric. Food Chem.*, 55: 7753-7757.